

Chemopreventive and antiproliferative effect of ANDROGRAPHIS PANICULATA EXTRACT

Agrawal RC^{1*}, Agrawal N¹, Mishra D¹

¹ Department of Research, Priyamvada Birla Cancer Research Institute, J. R. Birla Road, Satna-485005, Madhya Pradesh, India.

Correspondence: Agrawal RC. Email: rcagrawal60@yahoo.com

ABSTRACT

An *Andrographis paniculata* leaf and stem extract was studied in HeLa cells lines by In Vitro methods and anti promoting effect by skin tumour model. The dose dependent cytotoxicity was observed in HeLa cell lines by stem and leaves extracts of *Andrographis paniculata* extract. The prevention of bone marrow micronucleus formation by *Andrographis paniculata* leaves and stem extract was also observed. The reductions in tumour numbers were observed. The glutathione level was increased in the liver of animals which received the treatment of *Andrographis* extract along with DMBA + Croton Oil. The revealing information about the anticancer, antiproliferative and antimutagenic effect of an *Andrographis paniculata* extract was observed.

Key words: Micronucleus; Bone marrow; Glutathione; HeLa cells; Papilloma



ISSN:2397-9119

<http://mo.qingres.com>

OPEN ACCESS

DOI: 10.20900/mo.20170011

Received: March 10, 2017

Accepted: May 18, 2017

Published: June 25, 2017

Copyright: ©2017 Cain *et al.* This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

1 INTRODUCTION

Andrographis paniculata is used in Asia from centuries in traditional medicine to treat gastrointestinal (GI) tract and respiratory infectious diseases. It has been reported that *Andrographis* has a broad range of pharmacological effects^[1]. It has been suggested it's safe in controlled clinical trials report for treating upper respiratory tract infections. It also showed significant cardio protection by inducing antioxidant activity in myocardium^[2]. Cytotoxic activity against cancer cell lines has been reported by Compounds of *Andrographis paniculata*^[3]. Antimicrobial activity against eleven bacterial strains by ethanol extract of *Andrographis paniculata* have been reported^[4]. *Andrographolide* have been reported to hypoglycemic activity in rats^[5]. Antiulcer activity was reported in duodenal ulcer model in rats^[6]. Hepatoprotective effect was reported on acetaminophen induced hepatotoxicity in albino

rats^[7]. An andrographolide was also reported to induce apoptosis in TD-47 human breast cancer cell line in a time and concentration-dependent manner by increase expression of p53, bax, immuno histochemical parameters such as caspase-3 and decrease expression of bcl-2^[8]. *Andrographis paniculata*, dry leaf powder was reported to spermatogenesis, cessation of degenerative changes in the seminiferous tubules^[9]. The extract produced significant muscarinic activity, which accounts for its antivenom effects^[10]. Many of the conditions commonly treated with *Andrographis paniculata* in traditional medical systems are important, which requires further investigations for benefit in cancer treatment.

2 MATERIALS AND METHODS

2.1 Chemicals

The MTT cell culture tested reagent, 2,2 Diphenyl-1-picryl hydrazyl, 5,5 dithiobis (2 nitrobenzoic acid) were obtained from Hi media co., croton oil, 7,12 - Dimethylbenz (a) anthracene (DMBA), purchased from Sigma Chemicals Co. (St. Louis, MO. USA). Other chemicals were obtained Ranbaxy and SRL laboratories, Mumbai and purchased from local firms and were of the highest purity.

2.2 Animals

The experiment was approved by Institutional animal ethic committee before conduction of the experiments. These Swiss albino mice were housed in polypropylene cages at temperatures of 24 ± 30 .C. Mice were provided with standard mice feed and tap water ad libitum.

2.3 Preparation of *Andrographis paniculata* leaves and stem extract

Plant material (*Andrographis paniculata*) was collected and the specimen was authenticated by the botanist of Deen dayal Research Institute, Chitrakoot (MP), and India. Leaves and stem were washed, air dried, powdered and extracted separately using 50% methanol in a separating funnel. Extract thus obtained were vacuum evaporated into powder. These extract was again dissolved in double distilled water (DDW) immediately prior topical application. We have already done pharmacological screening of the components of extract in different solvent system such as methanol, Ethanol, Benzene and chloroform which showed the presence of carbohydrate,

flavonoid, Glycoside, Steroid, Resin, Tanin, Alkaloid.

2.4 Experimental design for Skin Carcinogenesis

One day before the experiment commenced, the dorsal skin on the animal's back was shaved. Two stage protocol was used as described by Berenblum^[11] and standardized by Agrawal *et al*^[12]. Animals were randomly allocated into 7 groups of comprising six mice each group. Single topical application of a carcinogen 7, 12-dimethylbenz (a) anthracene (DMBA) and then followed by a promoter (croton oil) twice in a week were used to induce tumours. The detail protocol was described in our earlier published paper^[17]. The treatment was provided topically to the shaved area. All animals groups were observed for gross and microscopic changes weekly All mice were weighed and examined for skin papillomas and results were recorded. The glutathione levels were measured by the method as described by Sedlak and Lindsay, 1968^[19].

2.5 MTT Assay

The extract of Stem and leaf of *Andrographis paniculata* on Hela cell line was studied. It was determined by MTT assay as described by Mosmann, 1983^[16]. For MTT assay, the extract at the dose of (0-1000 µg/ml and 5 µl/100 µg of cell suspension) was kept for 24 hours. Hela cells were cultured and treated with different concentrations of extract. 2 hours prior to the termination of experiment, MTT was added to cell culture at 0.25 mg/ml (5 µl of 5 mg/ml in 100 µl of cell suspension) concentration. At the end of the experiment, culture supernatant was removed and cell layer was dissolved in DMSO and further read in a plate reader at 550 nm and 660 nm. graph 1 and 2 plotted between the relative percentage death of cells against different concentration of *Andrographis* extract after 24 hour.

2.6 Micronucleus Assay

It was done by the method as described by Schmid (1975)^[13] and Aron *et al*^[14] and standardized by Agrawal *et al*^[15]. For the micronucleus assay, the extract at the volume of 0.2 ml at different doses level such as 500, 1000 and 1500 mg/kg body weight was injected 24 hours before the treatment of Cyclophosphamide to three animals. The positive control group received single ip. dose of Cyclophosphamide The cell were stained with May-Grunewald and Giemsa, and observed under microscope at 10X100X magnification. The detail protocol a described in our earlier paper Agrawal *et al*^[15].

3 RESULTS

The result showed that the animals of Group-V (control) produced skin papillomas, in all animals (100 %) which were observed from 5th week onwards. The significant tumour prevention was observed in the *Andrographis paniculata* extract treated experimental groups (25 % and 0 % in group VI & VII) compared to the carcinogen control. Cumulative number of papillomas was also reduced

in the *Andrographis paniculata* leaves and stem extract treated along with DMBA + Croton Oil groups (5 and 0 in group VI & VII). DMBA + Croton Oil treated showed Tumor burdens and tumor yields (3.5) where as it was decreased (0.5) in *andrographis* extract treated group (Table 1). The acute toxicity of *Andrographis* methanolic extract has been studied. It was observed that animals showed no toxicity up to the dose of 3000 mg/kg body weight in Swiss albino mice.

Table 1. Showing No. of Papilloma in the animals treated with *Andrographis paniculata* extract

Group	Treatment	Dose	Time of 1st appearance of Papilloma	Cumulative No. of Papilloma	Tumour yield	Tumour incidence
I	Vehicle alone	100µl/animal	–	–	–	–
II	DMBA alone	100µg/animal	–	–	–	–
III	Croton Oil alone	1% per animal	–	–	–	–
IV	<i>Andrographis</i> Extract alone	500mg/animal	–	–	–	–
V	DMBA + CO (Control)	100µg + 1% per animal	58th Day	21	21/6** (3.5)	6/6(100%)
VI	DMBA+ <i>Andrographis</i> (Leaves) Extract.+ CO	100µg + 1% + 500mg/animal	–	0	0/3	0/6 (0%)
VII	DMBA+ <i>Andrographis</i> (Stem)Extract.+ CO	100µg + 1% + 500mg/animal	61th Day	5	5/4	1/4* (25)

*Denotes statistically significant as compared to control in Students 't' test at $p < 0.05$. ** denotes total no. of Papilloma tumour/Total no. of animals

The induction of glutathione was also measured in the groups which were studied in papilloma models. The induction of glutathione level was significant in liver as compared to carcinogen control group. The result suggest significant increase

levels of Glutathione (GSH) in Liver of Papilloma bearing Swiss albino mice receiving treatment of *A. Paniculata* extract whereas in blood tissue it was almost similar to carcinogen control group and lower to untreated control group.


Table 2. Showing the level of Glutathione (GSH) in Blood and Liver


S.No	Groups	Treatment Group	Glutathione level	
			Blood (µg/ml)	Liver (µg/ml)
1	I	Normal mice	4.58±0.65	53.8±8.34
2	II	Carcinogen control (DMBA+CO)	2.7±0.06	14.95±9.75
3	III	DMBA+A.pan +CO(Leaves)	2.12±0.42	46.33±28.36*
4	IV	DMBA+A.pan +CO(Stem)	3.07±2.07	30.54±14.62*

*Denotes significant as compared to carcinogen control in students't' test. at $p < 0.05$.

The cytotoxicity of the plant was studied using HeLa cell lines in Vitro models. It shows that whole plant extract of *Andrographis paniculata*

had cytotoxic activity. Fig. 1 and 2 showed the relative percentage death of cells against different concentration of *Andrographis* extract after 24 hour.





In the micronucleus assay Cyclophosphamide (CP) has been used as a clastogen and anticlastogenic effect of *A. paniculata* (AP) has been observed in bone marrow cells of mice. The *Andrographis* extract showed a significant reduction in number of micronuclei as compared the CP alone. The result of Micronucleus assay showed that single application of *A. paniculata* hydro methanolic leaves

extract (i.p.) at the dose levels showed the significant reduction of micronucleus formation in PCE cell of bone marrow. However the micronucleus protection was not significant in stem extract of *Andrographis* extract treated group. The PCE/NCE ratio was comparable in all treated group as compared to CP which showed no toxicity of the extract in bone marrow cells of mice. (Table 3 & 4, Fig. 3)


Table 3. Effect of *A. paniculata* leaves extract on Micronucleus (MN) formation induced by Cyclophosphamide (CP) in bone marrow cells of Swiss albino mice.

S No.	Groups	Treatment Doses (mg/kg body wt)	MNPCE ± SEM	PCE/NCE Ratio ±SEM	% reduction in the frequency of CP induced DNA damage
1.	I	Cyclophosphamide Alone (50mg/kg b.wt)	3.5±1.1	0.99±0.08	-
2.	III	<i>A.paniculata</i> (500mg/kg b.wt)+ CP (50mg/kg b.wt)	2.2±1.3*	0.61±0.14	37.2%
3.	IV	<i>A.paniculata</i> (1000mg/kg b.wt)+ CP (50mg/kg b.wt)	1.6±0.8*	1.03±0.12*	55.3%
4.	V	<i>A.paniculata</i> (1500mg/kg b.wt)+ CP (50mg/kg b.wt)	0.5±0.3*	1.03±0.12*	85.8 %
5.	II	<i>A.paniculata</i> alone(1500mg/kg b.wt)	0.4 ±0.3	0.03 ±0.12	-
6.	VI	Vehicle alone (DDW)	0.16±0.16	0.44±0.08	-

*Denotes statistically significant as compared to cyclophosphamide group at $p < 0.05$.

Table 4. Effect of *A. paniculata* stem extract on Micronucleus (MN) formation induced by Cyclophosphamide (CP) in bone marrow cells of Swiss albino mice.

S No.	Groups	Treatment Doses (mg/kg body wt)	MNPCE±SEM	PCE/NCE Ratio ±SEM	% reduction in the frequency of CP induced DNA damage
1.	I	Cyclophosphamide Alone(50mg/kg b.wt)	3.5±1.1	0.99±0.08	-
2.	III	<i>A.paniculata</i> (500mg/kg b.wt)+ CP (50mg/kg b.wt)	2.5±1.22	0.62±0.20	28 %
3.	IV	<i>A.paniculata</i> (1000mg/kg b.wt)+ CP (50mg/kg b.wt)	2.6±0.89	0.587±0.12	25 %
4.	V	<i>A.paniculata</i> (1500mg/kg b.wt)+ CP (50mg/kg b.wt)	2.6±1.14	0.68±0.12	25 %
5.	II	<i>A.paniculata</i> alone(1500mg/kg b.wt)	0.82±0.66	0.53 ±0.054	-
6.	VI	Vehicle alone (DDW)	0.16.±0.16	0.44±0.08	-

**Fig. 3 Showing Micronucleus (MN) in polychromatic erythrocytes.**

4 DISCUSSIONS AND CONCLUSION

The current study demonstrates anticarcinogenic, antiproliferative and antimutagenic potential of *Andrographis paniculata* stem and leaves (AP) extract in skin papillomas, Hela cells lines and in bone marrow micronucleus assay. It demonstrates that a topical application of *andrographis paniculata* leaves and stem extract (500 mg/kg body weight) showed significant reduction in tumor incidence, in *andrographis* treated groups relative to the carcinogen treated control. In our previous study the anticarcinogenic activity of *andrographis paniculata* leaves has been reported^[17]. The methanolic extract of leaves of AP was more effective than stem extract in both Papilloma and micronucleus assay. The mechanism which shows induction of glutathione levels in liver tissues may be due to detoxification mechanism by liver of the animals. However the cytotoxicity anticarcinogenicity may be due to reactive oxygen species^[18]. The present study suggests the anticarcinogenic and antimutagenic activity of *Andrographis. paniculata* which is an important drug in traditional medicine and may be useful in treatment of cancer patients.

ACKNOWLEDGEMENTS

The authors are thanking to the M.P. Council of Science and Technology, Bhopal for its financial assistance and Dr. S.K. Maheshwari, Medical Director of PBCRI, Satna for providing facility to conduct the above work.

REFERENCES

- Sandberg F. *Andrographidis herba Chuanxinlian*: A review. Gothenburg, Sweden: Swedish Herbal Institute. Available from the American Botanical Council (USA). 1994.
- Ojha SK, Bharti S, Joshi S, Kumari S, Arya DS. Protective effect of hydroalcoholic extract of *Andrographis paniculata* on ischaemia-reperfusion induced myocardial injury in rats. *Indian J Med Res.* 2012; 135: 414-421.
- Aditya VS, Kumar NL, Mokkapati A. Evaluation of in vitro cytotoxicity of *Andrographis paniculata*, *Duranta serratifolia* and *Albizia lebeck* whole plant extracts by MTT assay against MCF-7 and HT-29 cell lines. *Cur Res Microbiol Biotechnol.* 2014; 2(2): 351-353.
- Mishra US, Mishra A, Kumari R, Murthy PN, Naik BS. Antibacterial activity of ethanol extract of *Andrographis paniculata*. *J Pharm Sci.* 2009; 71(4): 436-438.
- Nugroho AE, Andrie M, Warditiani NK, Siswanto E, Pramono S, Lukitaningsih E. Antidiabetic and antihyperlipidemic effect of *Andrographis paniculata* (Burm. f.) Nees and *andrographolide* in high-fructose-fat-fed rats. *Indian J Pharmacol.* 2012; 44(3): 377-381.
- Saranya P, Geetha A. Antiulcer activity of *Andrographis paniculata* (Burm.f.) Wall. against cysteamine-induced duodenal ulcer in rats. *Indian J Exp Biol.* 2011; 49(7): 525-533.
- Maleeka Begum, SF, Santhana Kumar, J and Swathi Balakrishnan, V, Antioxidant and hepatoprotective effect of *Andrographis paniculata* against acetaminophen (paracetamol) induced hepatotoxicity in Albino rats. *Int J Cur Res.* 2011;3(12):64-68
- Sukardiman H, Widyawaruyanti A, Sismindari, Zaini NC. Apoptosis Inducing Effect of *Andrographolide* on TD-47 Human Breast Cancer Cell Line. *African J Tradit Compliment Alternat Med.* 2007;4(3):345-351.
- Akbarsha MA, Murugaian P. Aspects of the male reproductive toxicity/male antifertility property of *andrographolide* in albino rats: effect on the testis and the cauda epididymidal spermatozoa. *Phytother Res.* 2000;14(6):432-435.
- Meenatchisundaram S, Parameswari G, Subbraj T, Suganya T, Michael A. Medicinal and Pharmacological Activities of *Andrographis paniculata* –Review. *Ethno botanical Leaflets.* 2009;13:55-58.
- Berenblum J, Shubik P. A new quantitative approach to the study of stages of chemical carcinogenesis in the mouse skin. *Br J Cancer.* 1947;1:373-391.
- Agrawal RC, Jain R, Raja W, Ovais M. Anticarcinogenic Effects of *Solanum lycopersicum* Fruit Extract on Swiss Albino and C57 BL Mice. *Asian Pac J Cancer Prev.* 2009;10:379-381.
- Schmid W. The micronucleus test. *Mutation Res.* 1975;31:9-15.
- Aron CS, Sorg R, Zimmer D. The mouse bone marrow micronucleus test. Evaluation of 21 drug candidates. *Mutation Res.* 1989;223:129-140.
- Agrawal RC, kumar S. Prevention of cyclophosphamide induced micronucleus formation in mouse bone marrow by indole-3 carbinol. *Food Clin Toxicol.* 1998;36:975-977.

16. Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods*. 1983;65(1-2):55-63.
17. Agrawal RC, Pandey P, Agrawal IN. Chemopreventive effects of *Andrographis paniculata* extract in Vivo and in Vitro models. *Med One*. 2016;1(4):1-6.
18. Ather M. Oxidative stress and experimental carcinogenesis. *Indian J Exp Biol*. 2002;40:656-667.
19. Sedlak J, Lindsay RH. Estimation of total protein-bound & non protein sulfhydryl group in tissues with Ellman's reagent. *Anal Biochem*. 1968;25:192-205.